March 2019

# HIP REPLACEMENTS – ORTHOPAEDICS – DR RIMMER – HE IS FAST!

Call patient 07:00am!

ALWAYS call – never ever wait for me - I know all the patients already! (E) As soon as we come in, call for the next patient (we call two ahead for arthroscopies) Have catheter setup ready (Surgical assistant (Grant) does unless I beat him to it) CONFIRM that blood is in Fridge (usually 2 units) – tell me it is in Fridge.

## Drugs:

**S8**: Midaz, Fentanyl Big 500mcg, Remifentanil 1mg, Suggamadex --- have ready Antibiotics: Vanc 1g, Ceph 2g, Gent 80mg

Others: Check drawer

Transexamic Acid 1g / Parecoxib 40mg / Tramadol 100mg Panadol 1g / Dex 8mg / Zofran 4mg / Maxalon 10mg Roc 50mg (a lot) / Ephedrine / Metaraminol (a lot) / Adrenaline (eek)

## Cannulation:

Please have a seat with wheels for me in anaesthetic bay Please have a desk for me in the operating room with another seat Cannula: 18g cannula / tegaderm / bung / Alco Wipe / Torniquet in a Kidney Dish If I miss twice I will ask for the ultrasound and frown.

# Monitoring & Table Supports:

- Try and have **two sets of ECG leads** (one for patient in bay, one for patient on table)
- When go into Theatre put **Sats on first** as soon as I see Sats > 95% I start giving drugs
- Connect drip set AFTER induction / Bear Hugger on / Don't bother Temp or BIS
- Side Arm boards, Upper arm support, Side Body Supports, & THREE **Towels**/patient.

#### Airways

- **ETT always** for hips (7.0 for F, 8.0 for M; I go down 0.5 size if I have trouble). Please have **KY Jelly** ready to put on tube ends, and make sure **tube cuff** is totally deflated, and disconnect the 10mL syringe from the end of the tube. Have a bougie ready.
- I use a C-Mac routinely (3 blade for F, 4 blade for M) for all my intubations on the Rimmer list – if I have trouble I then go to D-blade and definitely will need a bougie.
- I affix ETT with **brown tape** (Leukoplast not micropore): please have a pair of scissors available so I can make dog legs, along with an Alligator Extension (Expandable Tube Extension) and Tube Holder for once its all sorted.

# Intravenous & Pumps:

Pump Set with Extension and **Chook Foot** at the End – have some tape ready to affix. Hartmanns (warm if available) / Have a fluid warmer present and connected Two Syringe Drivers + Power Cables available (don't connect, I will trip over them).

Arterial Lines and Spinals and Post OT Blocks / Analgesia

I almost **never** do a spinal or arterial line for a Rimmer hip.

If I do an arterial line, ask me if I've booked a HDU Bed (I must be worried).

I will rarely do a Fascia Iliacus Block – 50mm stimuplex + 20mL 1% Naropin

I will always write an oxycodone PCA – please have a **PCA form** in theatre.